

# Generating velocity solutions with `globk`

M. A. Floyd   T. A. Herring

*Massachusetts Institute of Technology, Cambridge, MA, USA*

GNSS Data Processing and Analysis with GAMIT/GLOBK and `track`

UNAVCO Headquarters, Boulder, Colorado, USA

18–22 July 2022

[https://geoweb.mit.edu/gg/courses/202207\\_UNAVCO/](https://geoweb.mit.edu/gg/courses/202207_UNAVCO/)

Material from R. W. King, T. A. Herring, M. A. Floyd (MIT) and S. C. McClusky (now at ANU)

# Overview

- Basics of “velocity” solutions
  - Invoked with “apr\_neu all xx xx xx <NEU velocity sigmas>”
- Strategies for setting up solutions (they can take a long time to run)
- Strategies for speeding up solutions
- Methods for “cleaning up” potential problems
- Different reference frame realizations
- Some examples
- *These solutions involve making decisions about how to treat data and the type of solution to be created – lots of decisions*

# GLOBK velocity solutions

- The aim of these solutions is to combine many years of data to generate position, velocity, offset and postseismic parameter estimates
  - Increasingly common to have 10,000 parameters in these solutions if large networks over many years
- Input requirements for these solutions:
  - a priori coordinate and velocity file
    - Used as a check on positions in daily solutions (for editing of bad solutions) and adjustments are a priori values (a priori sigmas are for these values)
  - Earthquake file which specifies when earthquakes, discontinuities, and misnamed stations affect solution
    - Critical that this file correctly describe data.
  - Process noise parameters for each station
    - Critical for generating realistic standard deviations for the velocity estimates (e.g. `sh_gen_stats`).

# Velocity solution strategies

- In general careful setup (i.e. correct a priori coordinates, earthquake file and process noise files) is needed since each run that corrects a problem can take several days. Incorrect solutions may not complete correctly and results may be subtly wrong.
- General strategy for iteratively generating velocity solution:
  - Define a core-set of sites (usually 20-200 sites) where the solution runs quickly. Test files on this solutions and use the coordinate/velocity estimates to form the reference frame for time series generation.
  - Time series using these reference frame sites and then test (RMS scatter, discontinuity tests) to form a more complete earthquake and apriori coordinate/velocity files.
  - Steps above are repeated, usually increasing number of stations until solution is complete. As new stations are added missed discontinuities and bad process noise models can cause problems.
- Aim here is make sure that when a large solution is run (maybe several days of CPU time) that the run completes successfully.

# Before velocity runs

- Surveys may be combined into one solution per survey
- No need to re-run `glred` again to see long-term time series
- Multiple “.org”-files may be read by `tssum` or `sh_plot_pos`


```
tssum ts_pos mit.final_igb14 -R survey1_comb.org survey2_comb.org ...
```

- `ts_pos` is the name of a directory for the `.pos` files. (“.” can be used)


```
sh_plot_pos -f survey1_comb.org survey2_comb.org -k ...
```

# Example: Long-term time series for survey sites

## Reasonable repeatability



## Outlier in vertical



# Excluding outliers or segments of data

- Create “rename” file records and add to GLOBK command file’s “eq\_file” option, e.g.

```
rename PTRB      PTRB_XPS h1407080610_nb4a
rename PTRB      PTRB_XPS 2014 07 07 18 00 2014 07 08 18 30
rename ABCD      ABCD_XCL 2013 07 08 00 00
```

- “XPS” will not exclude data from `glred` (so still visible in time series) but will exclude data from `globk` (combination or velocity solution)
- “XCL” will exclude data from all `glred` or `globk` runs

# Run `globk`

- Create new “.gdl”-file with *combined* binary h-files, e.g. from `vsoln/`, assuming standard directory hierarchy

```
ls ../*/gsoln/*.GLX > vsoln.glx.gdl
```

- Optionally run `glist` to see size of solution
  - Recommended to prevent problems during long `globk` run
  - `glist` can read earthquake file and `globk` use site type commands (useful if a `globk` solution seems to be missing or has extra sites)
- Run `globk`
  - This may take many hours for very large/long velocity solutions
  - Use `tsfit` with earthquake file to generate a priori site coordinates. Be careful if `~/gg/tables/igb14_*.apr` files also used because some site names permutations may have inconsistent coordinates (use `unify_apr` to be safe)



# g\_lorg for different reference frames

- No need to re-run g\_lobk every time you want
- g\_lorg is usually called from g\_lobk command file (“org\_cmd” option) but g\_lorg may be run separately

```
globk 6 globk_vel.prt globk_vel.log globk_vel.gdl globk_vel.cmd  
glorg globk_vel_noam.org ERAS:... glorg_vel.cmd vel.com
```

- Must have saved the “.com”-file!
  - e.g. “com\_file @.com”
  - Do not use “del\_scra yes” in g\_lobk command file
  - “apr\_neu” must be loosely constrained (“apr\_rot” and “apr\_tran” will also need to be used for sestbl. “BASELINE” experiment solutions)

# Use of equates

- With earthquakes and discontinuities, there can be many site names for the same physically location:
  - Equate commands in `g1org` allow the velocity adjustments at these sites to be made the same (or constrained to be the same within a specified sigma)
  - “`eq_dist`” allows site separate by distance to equated (and constrained in latest `g1org`)
  - “`eq_4char`” equates sites with same 4-character name (useful to stop equates at sites that share antennas)
  - chi-squared increments of equates allows assessment of equates (use “`unequate`” for large chi-squared values)
  - Use “`FIXA`” option to make a priori the same for equated sites (better to use consistent a priori file)

# Uses of `sh_gen_stats`

- Velocity solutions are often iterative:
  - Generate time series using some reference frame sites (IGb14 sites initially for example)
  - Fit to the time series (`tsfit`) to:
    - Find outliers, nature of earthquakes (log needed?), discontinuities
    - Self consistent a priori file.
    - Used FOGMEx model (realistic sigma) to get process noise model and list of low-correlated noise reference frame sites). Use “stabrad” option for dense networks
  - Run `g1obk` velocity solution to refine reference frame site coordinates and velocities
  - Re-generate time series and repeat

# Some comparisons: Approach

- Use `sh_exglk -f <soln.org> -vel <soln.vel> -rmdup` to extract velocity estimates (rmdup removes equated sites with the same estimates)
- Program `velrot` allows fields to be compared (change frames and merge fields as well), for example:  
`velrot solna.vel nam14 solnb.vel IGb14 ' ' ' ' N`  
compares to solutions directly (use “RT” instead of “N” to allow rotation and translation rates)
  - Use “`grep '^S '`” to get statistics

# Comparisons: Decimation

- Decimation: Different days of week (1996-2015 solution, small subset of sites):

Un-aligned fields

```
compare 1 NMT_vel_150418_day1.vel NMT_vel_150418_day3.vel
S Component North      #      75 WMean  -0.00 WRMS   0.04 mm/yr, NRMS   0.198
S Component East       #      75 WMean  -0.02 WRMS   0.04 mm/yr, NRMS   0.203
S Component Up         #      75 WMean   0.03 WRMS   0.16 mm/yr, NRMS   0.180
S Component Horz      #      75 WMean  -0.01 WRMS   0.04 mm/yr, NRMS   0.200
compare 2 NMT_vel_150418_day1.vel NMT_vel_150418_day5.vel
S Component North      #      74 WMean  -0.01 WRMS   0.04 mm/yr, NRMS   0.207
S Component East       #      74 WMean  -0.02 WRMS   0.05 mm/yr, NRMS   0.225
S Component Up         #      74 WMean   0.04 WRMS   0.19 mm/yr, NRMS   0.212
S Component Horz      #      74 WMean  -0.01 WRMS   0.04 mm/yr, NRMS   0.217
compare 3 NMT_vel_150418_day3.vel NMT_vel_150418_day5.vel
S Component North      #      76 WMean  -0.01 WRMS   0.03 mm/yr, NRMS   0.177
S Component East       #      76 WMean  -0.01 WRMS   0.03 mm/yr, NRMS   0.161
S Component Up         #      76 WMean   0.01 WRMS   0.13 mm/yr, NRMS   0.142
S Component Horz      #      76 WMean  -0.01 WRMS   0.03 mm/yr, NRMS   0.169
```

# Comparison: Time series vs GLOBK

- PBO Combined analyses:

Un-aligned fields (no rotation and translation).

compare 1 PBO\_vel\_150425.vel PBO\_vel\_150425KF.vel

S Component North	#	2105	WMean	-0.01	WRMS	0.12	mm/yr,	NRMS	0.925
S Component East	#	2105	WMean	-0.00	WRMS	0.13	mm/yr,	NRMS	0.934
S Component Up	#	2105	WMean	0.02	WRMS	0.31	mm/yr,	NRMS	0.871
S Component Horz	#	2105	WMean	-0.01	WRMS	0.12	mm/yr,	NRMS	0.929

compare 4 PBO\_vel\_150425.vel PBO\_vel\_150425\_NAM08.vel

S Component North	#	1972	WMean	0.03	WRMS	0.13	mm/yr,	NRMS	0.965
S Component East	#	1972	WMean	0.02	WRMS	0.15	mm/yr,	NRMS	1.049
S Component Up	#	1972	WMean	-0.07	WRMS	0.41	mm/yr,	NRMS	0.943
S Component Horz	#	1972	WMean	0.02	WRMS	0.14	mm/yr,	NRMS	1.008

compare 7 PBO\_vel\_150425KF.vel PBO\_vel\_150425\_NAM08.vel

S Component North	#	1969	WMean	0.04	WRMS	0.16	mm/yr,	NRMS	0.952
S Component East	#	1969	WMean	0.02	WRMS	0.17	mm/yr,	NRMS	0.967
S Component Up	#	1969	WMean	-0.08	WRMS	0.44	mm/yr,	NRMS	0.935
S Component Horz	#	1969	WMean	0.03	WRMS	0.16	mm/yr,	NRMS	0.959

PBO\_vel\_150425.vel: tsfit solution to time series

PBO\_vel\_150425KF.vel: tsfit Kalman filter solution to timeseries

PBO\_vel\_150425\_NAM08.vel: GLOBK combined velocity solution (NMT+CWU), decimated 7 days, 28-subnet combination.

Reference frame realization to NAM08 frame sites (~600)

- See Herring et al., Reviews of Geophysics, 2016 for more detailed comparisons

# Final comments

- Practice large solutions with decimated data sets and small networks (run time increased cubically with number of stations)
- Make sure your a priori coordinates files are consistent (especially with equates)
  - Use the `out_aprf` command in `tsfit` to generate an a priori which is consistent with your timeseries estimates